

AGENT FOR PREVENTING RECURRENCE OF CEREBROVASCULAR DISORDER  
AND AGENT FOR AMELIORATING TROUBLES FOLLOWING  
CEREBROVASCULAR DISORDER AND INHIBITING PROGRESS THEREOF

5 Technical Field

10 The present invention relates to an agent for preventing recurrence of cerebrovascular disorder comprising a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof as an active ingredient, as well as an agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof comprising a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof as an active ingredient.

15 Background Art

20 According to the classification of cerebrovascular disorder, 3rd edition (MINDS-III, Stroke 21:637-676, 1990), National Institute of Neurological Disorders and Stroke (MINDS), cerebrovascular disorder is classified into asymptomatic cerebral infarction, transient ischemic attack (TIA), cerebral apoplexy, cerebrovascular dementia, and hypertensive encephalopathy. The type of cerebral apoplexy includes cerebral hemorrhage, subarachnoid hemorrhage, 25 cranial hemorrhage accompanying a malformation in cerebral

arteries and veins, and cerebral infarction, and there are many diseases causing each of them. In the past, hypertensive cerebral hemorrhage was a major disease, but by finding and managing hypertension at an early stage and westernization of nutrition intake, the rate of occurrence thereof and the death rate are gradually reduced, while the absolute number of old men with cerebral infarction and the ratio thereof are increasing due to spread of various kinds of image diagnosis techniques. Cerebral infarction is classified into atherothrombotic cerebral infarction, lacuna infarction, and cardiogenic cerebral infarction, among which lacuna infarction is decreasing with the progress of treatment of hypertension. On the other hand, the ratio of atherothrombotic cerebral infarction and cardiogenic cerebral infarction is increasing due to westernization of nutrition intake.

In Japan, these cerebrovascular disorders are at the second place of causes of death or at the top of causative diseases for bedridden old men at present, and thus countermeasures not only for prevention of occurrence of cerebrovascular disorder but also for prevention of recurrence thereof and progress of troubles following cerebrovascular disorder become important problems.

Medical treatment of cerebrovascular disorder is divided by a stage of diseases (acute stage, subacute stage,

10031398-011802

chronic stage, asymptomatic stage), a type of diseases (hemorrhagic, ischemic, etc.) and severity of diseases (presence or absence of hernia).

5 Treatment of cerebral apoplexy at an acute stage and a subacute stage includes systemic management including blood pressure, as well as inhibition of cerebral edema, measures against complications (hemorrhage in digestive tracts, infections in respiratory organs, convulsions, unrest, etc.), brain-protecting therapy (treatment by a brain-protecting drug, cryotherapy, etc.) and antithrombotic therapy depending on a particular type of diseases (treatment with thrombolytics, antiplatelets, anticoagulants, etc.), etc. For cerebral apoplexy at an acute stage, it is necessary to take a measure against cerebral edema in every type of diseases. Cerebral edema occurring at an acute stage of cerebrovascular disorder is a phenomenon in which the blood-brain barrier is disturbed by damage to brain, and fluid components in brain tissues are increased to increase the volume of brain, against which chemotherapy is generally conducted by intermittent administration of steroids or by intravenous injection of a hypertonic solution such as mannitol or glycerin. At an acute stage of cerebral apoplexy, in addition to a reactive increase in blood pressure, there are various changes in cerebral circulation kinetics, that is, a decrease in a

10  
15  
20  
25

10031398 011802  
202510 081100T

cerebral blood flow rate not only in foci but also in a surrounding penumbra region and in a part distant from foci in the brain, disorder in the ability for automatic regulation of cerebral blood flow, a reduction in CO<sub>2</sub> reactivity in cerebral vessels, etc. Accordingly, there is danger that a careless and significant reduction in blood pressure permits cerebral blood flow to be reduced depending on blood pressure and a penumbra region can be exposed to irreversible danger. Then, use of an antihypertensive is restricted in principle, unless complications with severe hypertension and organ disorders attributable to hypertension such as heart failure are present. In the case of cerebral hemorrhage, however, there is danger of increasing hematocoele, and thus about 20% reduction in blood pressure is often recommended, and in the case of subarachnoid hemorrhage, an increase in blood pressure causes re-bleeding and significantly influences the prognosis of life, and thus a durable depression in blood pressure is required.

Major medical treatment at a chronic stage involves a measure against troubles following cerebrovascular disorder and a measure for preventing recurrence thereof, depending on a particular type of diseases. The troubles following cerebrovascular disorder are classified roughly into nerve symptoms, mind symptoms, subjective symptoms (non-

localizing subjective symptoms, localizing subjective symptoms, etc.), and obstacles in activities of daily living (ADL). In chemotherapy for these troubles following cerebrovascular disorder, nerve symptoms are limited to symptomatic therapy by an antispasmodic, etc. Chemotherapy is mainly directed to amelioration of nerve symptoms and subjective symptoms and uses agents for ameliorating cerebral circulation, agents for ameliorating cerebral metabolism (neurotransmission ameliorating agents, cerebral metabolism-activating agents, etc.) and psychotropics (anti-schizophrenia, anti-anxiety, antidepressant, etc.). Major treatment of obstacles in ADL involves rehabilitation, and chemotherapy is used as an aid to promote rehabilitation. Measures for preventing recurrence of cerebrovascular disorder include not only treatment (indirect measures) of diseases serving as factors causing hypertension, diabetes, hyperlipemia, etc., but also chemotherapy using antiplatelets, anti-coagulants, etc. depending on a particular type of diseases. However, there is no report suggesting that a compound having an AII antagonistic activity can be used in a direct measure for preventing recurrence of cerebrovascular disorder or for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof.

While treatment at an acute stage is the most

10031398-011802

important for treatment of cerebrovascular disorder from the viewpoint of medical economy, prevention of recurrence thereof and its progress are also major problems. However, only an antiplatelet is used for preventing recurrence of cerebrovascular disorder, and its usefulness is low.

The most dangerous factor for cerebrovascular disorder is hypertension, while the dangerous factors for cerebral infarction include an abnormality of sugar resistance and an abnormality in electrocardiogram, and the dangerous factors for cerebral hemorrhage include an abnormality in electrocardiogram, an abnormality in eye ground and drinking, and the like. As the dangerous factors for recurrence of cerebrovascular disorder, hypertension, cardiac disorders, TIA, diabetes, etc. are pointed out, and it can be considered that antihypertensive therapy is applied not only to prevention of occurrence of cerebrovascular disorder (primary prevention) but also to prevention of recurrence thereof (secondary prevention). However, usually hypertension accompanying cerebral apoplexy causes complications with diabetes, hyperlipemia and obesity, and accompanies high rates of vascular lesions in hearts, kidneys and peripheral arteries, and in such conditions, a presently existing antihypertensive therapy does not necessarily prevent the recurrence of cerebral apoplexy. This is because, in cerebral infarction after

10031398 011802

the onset, it is considered that cerebral arteriosclerotic lesions also proceed, and conventional antihypertensives cannot be expected to ameliorate such vascular lesions. Further, atherothrombotic cerebral infarction accompanying  
5 constriction and occlusion of a major stem artery occurs due to kinetics of the blood flow mechanism, so hypotension is considered to cause cerebral ischemia. In particular, the ability for automatic regulation of cerebral circulation has failed at an acute stage of cerebral  
10 apoplexy, and therefore rapid depression in blood pressure could conversely cause a reduction in cerebral blood flow to permit progress of cerebral ischemia. Further, at a chronic stage of 1 month or more after the onset of disease conditions, wherein antihypertensive therapy is usually  
15 initiated, the upper and lower limits of automatic regulation ability are deviated rightward (toward higher blood pressure level) in old men or patients with hypertension and cerebrovascular disorder, so even a slight depression in blood pressure causes a reduction in cerebral  
20 blood flow, to cause recurrence of cerebral ischemia. Further, patients with hypertension and cerebrovascular disorder often exhibit a non-dipper where blood pressure at night is increased even by administration of an antihypertensive, or an extra-dipper where blood pressure  
25 at night is too lowered, and hypertension is caused in

10031398-011802

early morning, and this change in blood pressure serves as a cause for inducing recurrence of cerebral apoplexy. Accordingly, there is demand for a drug having an anti-arteriosclerosis action, not lowering blood flow, improving the ability for automatic regulation of cerebral blood flow, and showing a stable and durable antihypertensive action.

At present, a long-lasting Ca antagonist or an angiotensin converting enzyme (ACE) inhibitor is used as an antihypertensive as a first choice in consideration of its action on cerebral circulation and cerebral vessels, and other drugs used include vasodilative  $\beta$ - or  $\alpha_1$ -blockers. However, these drugs have problems with significant depression in blood pressure, duration of their action, and their side effects, and at present there is no ideal drug. Among them, due to a mechanism which is presumed to be attributable to a reduction in angiotensin II (AII) production in cerebral endarterium, the ACE inhibitor can particularly deviate the ability for automatic regulation of cerebral circulation leftward (toward lower blood pressure level), thus preventing or reducing cerebral ischemia developed by kinetics of the blood flow mechanism, and in this respect, the ACE inhibitor is a useful drug. However, the ACE inhibitor has the activity of not only inhibiting AII production but also decomposing inflammatory mediators such as bradykinin and substance P, and because



of coughs and vascular edemas resulting therefrom,  
administration of the drug should inevitably be terminated  
in many cases. Further, an enlargement of edemas and  
cellular disorder may be induced due to enhancement of the  
inflammatory action, and particularly pathema at an acute  
stage (including ultra-acute stage to subacute stage)  
significantly influencing prognosis could be significantly  
worsened. Further, at present, there are little drugs  
(excellent in T/P ratio) showing stable depression in blood  
pressure through 24 hours, and in patients exhibiting a  
non-dipper having recurrence of cerebrovascular disorder at  
high rate or patients exhibiting hypertension in early  
morning, sufficiently stable depression in blood pressure  
cannot be achieved.

On the other hand, it is described that a compound  
having an antagonism to AII is known as an agent for  
treating diseases in circulatory organs, such as  
hypertension, cardiac diseases (cardiac hypertrophy, heart  
failure, myocardial infarction, etc.), cerebral apoplexy,  
nephritis, etc. (JP-A 4-364171, etc.), and it is  
demonstrated that AII having a strong vasoconstrictor  
action is prevented from acting on AII receptors by this  
compound to exhibit a durable action of depressing blood  
pressure.

Further, it is reported that Candesartan, i.e. an

active metabolite of Candesartan cilexetil can deviate the lower limit of automatic regulation leftward (toward lower blood pressure level) (Vraamark T et al., J Hypertens 13:755-761, 1995), but there is no report suggesting that a compound having AII antagonistic activity is useful as an agent for preventing recurrence of cerebrovascular disorder or as an agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof.

10 Disclosure of the Invention

The present invention provides pharmaceutical preparations useful for preventing recurrence of cerebrovascular disorder and for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof. Cerebrovascular disorder causes various troubles following the onset thereof, and therefore reduction of troubles following the cerebrovascular disorder and prevention the recurrence thereof are not only very important for a patient's comeback to normal life, but also can significantly contribute to a reduction in the number of bedridden old men or patients with vascular dementia, to significantly reduce medical expenses and the burden on the family.

Under the above-mentioned circumstances, the present inventors intensively studied pharmaceutical preparations

20031398.01802

useful for preventing recurrence of cerebrovascular disorder and for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof, and as a result, they found that a compound having an angiotensin II antagonistic activity, especially a compound represented by a specific structural formula and having angiotensin II (AII) antagonistic activity is very effective in preventing recurrence of cerebrovascular disorder and in ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof, and as a result of further examination on the basis of these findings, the invention was completed.


That is, the present invention relates to:

- (1) an agent for preventing recurrence of cerebrovascular disorder or an agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof which comprises a compound having an angiotensin II antagonistic activity (that is, a compound having an angiotensin II receptor antagonistic activity), a prodrug thereof or a salt thereof;
- (2) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity is a non-peptide compound;
- (3) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity

is a compound having an oxygen atom in the molecule;

(4) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity is a compound having an ether bond or a carbonyl group:

5 (5) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity is a compound represented by the formula (I):



wherein  $R^1$  represents a group capable of forming an anion or being converted into said group, X indicates that the phenylene group and the phenyl group are bound to each other directly or via a spacer of a chain made of 2 or less atoms, n is an integer of 1 or 2, ring A represents a benzene ring which may further have substituent(s),  $R^2$  represents a group capable of forming an anion or being converted into said group, and  $R^3$  represents a hydrocarbon residue which may be bound via a heteroatom and may have substituent(s);

(6) the agent according to the above-mentioned (1), wherein

the compound having an angiotensin II antagonistic activity is Losartan, Eprosartan, Candesartan cilexetil, Candesartan, Valsartan, Telmisartan, Irbesartan, Olmesartan or Tasosartan;

5 (7) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity is 2-ethoxy-1-[[2'-(1H-tetrazole-5-yl)biphenyl-4-yl]methyl]benzimidazole-7-carboxylic acid;

10 (8) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity is 1-(cyclohexyloxycarbonyloxy)ethyl 2-ethoxy-1-[[2'-(1H-tetrazole-5-yl)biphenyl-4-yl]methyl]benzimidazole-7-carboxylate;

15 (9) the agent according to the above-mentioned (1), wherein the compound having an angiotensin II antagonistic activity is 2-ethoxy-1-[[2'-(2,5-dihydro-5-oxo-1,2,4-oxadiazole-3-yl)biphenyl-4-yl]methyl]benzimidazole-7-carboxylic acid;

(10) the agent according to the above-mentioned (1), wherein the troubles following cerebrovascular disorder are nerve symptoms;

(11) the agent according to the above-mentioned (1), wherein the troubles following cerebrovascular disorder are mental symptoms;

25 (12) the agent according to the above-mentioned (1), wherein the troubles following cerebrovascular disorder are

10034393 "01B02"

subjective symptoms;

(13) the agent according to the above-mentioned (1), wherein the troubles following cerebrovascular disorder are obstacles in activities of daily living;

5 (14) use of a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof for preventing recurrence of cerebrovascular disorder or in ameliorating troubles following cerebrovascular disorder or inhibiting progress thereof;

10 (15) use of a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof for manufacturing a pharmaceutical preparation for preventing recurrence of cerebrovascular disorder or for ameliorating troubles following cerebrovascular disorder or  
15 inhibiting progress thereof;

(16) a method for preventing recurrence of cerebrovascular disorder in mammals which comprises administering an effective amount of a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof  
20 to the mammals;

(17) a method for ameliorating troubles following cerebrovascular disorder or inhibiting progress thereof in mammals which comprises administering an effective amount of a compound having an angiotensin II antagonistic  
25 activity, a prodrug thereof or a salt thereof to the

10031393 "011302"

mammals;

- (18) use of a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof for manufacturing a pharmaceutical preparation for preventing recurrence of cerebrovascular disorder; and
- (19) use of a compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof for manufacturing a pharmaceutical preparation for ameliorating troubles following cerebrovascular disorder or inhibiting progress thereof.

In the present invention, the compound having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof can be advantageously used for preventing recurrence of cerebrovascular disorder or ameliorating troubles following cerebrovascular disorder or inhibiting progress thereof, and for preventing occurrence of asymptomatic cerebral infarction or inhibiting transition thereof into symptomatic cerebral infarction, or for inhibiting transition of troubles following cerebrovascular disorder into cerebrovascular dementia.

The more specific action of the compound of the present invention having an angiotensin II antagonistic activity, a prodrug thereof or a salt thereof includes, for example, inhibition or amelioration of cerebral edema after cerebrovascular disorder; inhibition or amelioration of

cellular disorder after cerebrovascular disorder;  
inhibition or amelioration of cerebral neuron disorder  
after cerebrovascular disorder; protection of brain after  
cerebrovascular disorder; inhibition or amelioration of  
5 endothelial cellular disorder after cerebrovascular  
disorder; inhibition or amelioration of infiltration of  
inflammatory cells or inflammation after cerebrovascular  
disorder; inhibition of promotion of free radical  
production from inflammatory cells, etc. after  
10 cerebrovascular disorder; inhibition of production and/or  
action of vascular agonists and thrombus inducers whose  
production increases after cerebrovascular disorder;  
amelioration of microcirculation after cerebrovascular  
disorder; inhibition of a promoting action on vascular  
15 penetration in cerebrovascular disorder; amelioration of  
disorder in the penumbra region after cerebrovascular  
disorder; amelioration of the cerebral blood barrier after  
cerebrovascular disorder; inhibition of an increase in  
hematocrit (increase in blood viscosity) after  
20 cerebrovascular disorder; inhibition of formation of  
thrombi and emboli causing reoccurrence of cerebrovascular  
disorder; amelioration of vascular endothelial cell  
disorder causing reoccurrence of cerebrovascular disorder;  
amelioration of cardiac function disorder causing  
25 reoccurrence of cerebrovascular disorder, particularly

10034398-011802



cardiogenic cerebral infarction; amelioration of an  
abnormality in the transformability of erythrocytes after  
cerebrovascular disorder; inhibition of promotion of the  
agglutination ability of erythrocytes after cerebrovascular  
5 disorder; inhibition of promotion of the agglutination  
ability of platelets after cerebrovascular disorder;  
inhibition of promotion of the adherence ability of  
leukocytes after cerebrovascular disorder; inhibition or  
amelioration of the progress of arteriosclerosis foci in  
10 vascular lesions causing recurrence of cerebrovascular  
disorder, particularly cerebral infarction; etc.

In the present invention, the term "angiotensin II  
antagonistic activity" refers to the action of  
competitively or noncompetitively inhibiting binding of  
15 angiotensin II to angiotensin II receptors on cell  
membranes to attenuate strong vasoconstricting action and  
vascular smooth muscle proliferating action induced by  
angiotensin II to mitigate symptoms of hypertension.


The compound having an angiotensin II antagonistic  
20 activity used in the present invention may be a peptide or  
non-peptide compound, preferably a non-peptide antagonistic  
compound because of the advantage of long-lasting action.  
The compound having an angiotensin II antagonistic activity  
is preferably a compound having an oxygen atom in the  
25 molecule, particularly preferably a compound having an

ether bond or a carbonyl group (said carbonyl group may form a hydroxyl group due to resonance), more preferably a compound having an ether bond or a ketone derivative, most preferably an ether derivative.

5           The non-peptide compounds having an angiotensin II antagonistic activity are not particularly limited, but, for example, imidazole derivatives are disclosed in JP-A 56-71073, JP-A 56-71074, JP-A 57-98270, JP-A 58-157768, USP 4,355,040 and USP 4,340,598; improved imidazole derivatives  
10       are disclosed in EP-253310, EP-291969, EP-324377, EP-403158, WO-9100277, JP-A 63-23868 and JP-A 1-117876; pyrrole, pyrazol and triazole derivatives are disclosed in USP 5,183,899, EP-323841, EP-409332 and JP-A 1-287071; benzimidazole derivatives are disclosed in USP 4,880,804,  
15       EP-0392317, EP-0399732, EP-0400835, EP-425921, EP-459136 and JP-A 3-63264; azaindene derivatives are disclosed in EP-399731 etc.; pyrimidone derivatives are disclosed in EP-407342 etc.; quinazoline derivatives are disclosed in EP-411766 etc.; xanthine derivatives are disclosed in EP-  
20       430300 etc.; condensed imidazole derivatives are disclosed in EP-434038 etc.; pyrimidine dione derivatives are disclosed in EP-442473 etc.; thienopyridone derivatives are disclosed in EP-443568 etc.; and heterocyclic compounds are disclosed in EP-445811, EP-483683, EP-518033, EP-520423,  
25       EP-588299 and EP-603712. Among these, representative

compounds are also described in Journal of Medicinal Chemistry (Vol. 39, No. 3, pp. 625-656, 1996). As the non-peptide compounds having an angiotensin II antagonistic activity, it is possible to use, in addition to the compounds described above in the prior art literatures, any non-peptide compounds having an angiotensin II antagonistic activity, among which preferably used are Losartan (DuP753), Eprosartan (SK&F108566), Candesartan cilexetil (TCV-116), Valsartan (CGP-48933), Telmisartan (BIBR277), Irbesartan (SR47436) and Tasosartan (ANA-756), Olmesartan (CS-866), as well as active metabolites thereof (Candesartan, etc.), etc.

As the non-peptide compounds having an angiotensin II antagonistic activity, it is also preferable to employ, for example, a benzimidazole derivative represented by the formula (I):





wherein  $R^1$  represents a group capable of forming an anion or being converted into said group, X indicates that the phenylene group and the phenyl group are bound to each

other directly or via a spacer of a chain made of 2 or less atoms, n is an integer of 1 or 2, ring A represents a benzene ring which may further have substituent(s),  $R^2$  represents a group capable of forming an anion or being converted into said group, and  $R^3$  represents a hydrocarbon residue which may be bound via a heteroatom and may have substituent(s) (preferably a hydrocarbon residue which is bound via an oxygen atom and may have substituent(s)) or a salt thereof.

As  $R^1$  in the above-mentioned formula (I), the group capable of forming an anion (that is, a group having a hydrogen atom releasable as a proton) includes, for example, (1) a carboxyl group, (2) a tetrazolyl group, (3) a trifluoromethanesulfonic acid amide group ( $-NHSO_2CF_3$ ), (4) a phosphate group, (5) a sulfonate group, (6) a 5- to 7-membered (preferably 5- to 6-membered) monocyclic, optionally substituted heterocyclic residue containing at least one atom selected from N, S and O.


Examples of the "5- to 7-membered (preferably 5- to 6-membered) monocyclic, optionally substituted heterocyclic residue containing at least one atom selected from N, S and O" include:






and the like.

- 5 In the formulas above, when g represents -NH- etc., the bond between the heterocyclic residue represented by  $R^1$  and the phenyl group to which the heterocyclic residue is bound may be not only the carbon-carbon bond, but also one of plural nitrogen atoms present therein. For example, if
- 10  $R^1$  is represented by:





then, the bond therebetween is shown by, for example:



Other examples of such bonds via a nitrogen atom

5 include:




and the like.

In the formulas above, g represents  $-\text{CH}_2-$ ,  $-\text{NH}-$ ,  $-\text{O}-$   
 10 or  $-\text{S}(\text{O})_m-$ ;  $>=Z$ ,  $>=Z'$  and  $>=Z''$  represent a carbonyl group,

a thiocarbonyl group and an optionally oxidized sulfur atom (e.g., S, S(O), S(O)<sub>2</sub> etc.) (preferably a carbonyl or thiocarbonyl group, more preferably a carbonyl group) respectively; and m is an integer of 0, 1 or 2.

5            Preferably, the heterocyclic residue represented by R<sup>1</sup> is, for example, a group having both an -NH- or -OH- group as a proton donor such as in an oxadiazolone ring, an oxadiazolothion ring or a thiadiazolone ring and a carbonyl group, a thiocarbonyl group or a sulfinyl group as a proton  
10            acceptor. Further, the heterocyclic residue represented by R<sup>1</sup> may form a condensed ring by binding a cyclic substituent thereto, but the heterocyclic residue represented by R<sup>1</sup> is preferably a 5- to 6-membered ring, more preferably a 5-membered ring.

15            The heterocyclic ring represented by R<sup>1</sup> is preferably a group represented by the formula:




wherein i represents -O- or -S-, j represents >=O, >=S or >=S(O)<sub>m</sub> wherein m has the same meanings as defined above,  
20            and said heterocyclic ring is particularly preferably 2,5-dihydro-5-oxo-1,2,4-oxadiazole-3-yl, 2,5-dihydro-5-thioxo-1,2,4-oxadiazole-3-yl, 2,5-dihydro-5-oxo-1,2,4-thiadiazole-




3-yl, especially 2,5-dihydro-5-oxo-1,2,4-oxadiazole-3-yl.


As the above heterocyclic residue ( $R^1$ ), tautomers are present as shown below. For example, in the case of:



- 5 wherein  $Z = O$  and  $g = O$ , there occur three  $a'$ ,  $b'$  and  $c'$  tautomers as shown below:



and the heterocyclic residue represented by:



- 10 includes all the  $a'$ ,  $b'$  and  $c'$  mentioned above.

As  $R^1$ , the group capable of forming an anion may have

been protected at replaceable positions by an optionally substituted lower ( $C_{1-4}$ ) alkyl group or an acyl group (e.g., lower ( $C_{2-5}$ ) alkanoyl, benzoyl etc.).

The optionally substituted lower ( $C_{1-4}$ ) alkyl group includes, for example, (1) a lower ( $C_{1-4}$ ) alkyl group which may be substituted with 1 to 3 phenyl groups which may have a halogen atom, nitro, lower ( $C_{1-4}$ ) alkyl, lower ( $C_{1-4}$ ) alkoxy etc. (e.g., methyl, triphenylmethyl, p-methoxybenzyl, p-nitrobenzyl etc.), (2) a lower ( $C_{1-4}$ ) alkoxy-lower ( $C_{1-4}$ ) alkyl group (e.g., methoxymethyl, ethoxymethyl etc.), and (3) the formula  $-CH(R^4)-OCOR^5$  wherein  $R^4$  represents (a) hydrogen, (b) a  $C_{1-6}$  straight or branched lower alkyl group (e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, isopentyl, neopentyl etc.), (c) a  $C_{2-6}$  straight or branched lower alkenyl group or (d) a  $C_{3-8}$  cycloalkyl group (e.g., cyclopentyl, cyclohexyl, cycloheptyl etc.), and  $R^5$  represents (a) a  $C_{1-6}$  straight or branched lower alkyl group (e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl etc.), (b) a  $C_{2-6}$  straight or branched lower alkenyl group, (c) a  $C_{1-3}$  lower alkyl group which may be substituted with a  $C_{3-8}$  cycloalkyl group (e.g., cyclopentyl, cyclohexyl, cycloheptyl etc.) or with an aryl group which may be substituted (e.g., a phenyl or naphthyl group which may have a halogen atom, nitro, lower ( $C_{1-4}$ )

10034398-011802

alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) (e.g., benzyl, p-chlorobenzyl, phenethyl, cyclopentylmethyl, cyclohexylmethyl etc.), (d) a C<sub>2-3</sub> lower alkenyl group substituted with C<sub>3-8</sub> cycloalkyl or with an aryl group which may be substituted (e.g., a phenyl or naphthyl group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) (e.g., a group such as cinnamyl having an alkenyl moiety such as vinyl, propenyl, allyl, isopropenyl etc.), (e) an aryl group which may be substituted (e.g., a phenyl or naphthyl group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc., such as phenyl, p-tolyl, naphthyl etc.), (f) a C<sub>1-6</sub> straight or branched lower alkoxy group (e.g., methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec-butoxy, t-butoxy, n-pentyloxy, isopentyloxy, neopentyloxy etc.), (g) a C<sub>2-8</sub> straight or branched lower alkenyloxy group (e.g., allyloxy, isobutenyloxy etc.), (h) a C<sub>3-8</sub> cycloalkyloxy group (e.g., cyclopentyloxy, cyclohexyloxy, cycloheptyloxy etc.), (i) a C<sub>1-3</sub> lower alkoxy group substituted with a C<sub>3-8</sub> cycloalkyl (e.g., cyclopentyl, cyclohexyl, cycloheptyl etc.) or with an aryl group which may be substituted (e.g., a phenyl or naphthyl group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) (e.g., a group such as benzyloxy, phenethyloxy, cyclopentylmethoxy, cyclohexyl methoxy etc.

20031398-011802

having an alkoxy moiety such as methoxy, ethoxy, n-propoxy, isopropoxy etc.), (j) a C<sub>2-3</sub> lower alkenyloxy group substituted with a C<sub>3-8</sub> cycloalkyl (e.g., cyclopentyl, cyclohexyl, cycloheptyl etc.) or with an aryl group which may be substituted (e.g., a phenyl or naphthyl group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) (e.g., a group such as cinnamyloxy etc. having an alkenyloxy moiety such as vinyloxy, propenyloxy, allyloxy, isopropenyloxy etc.), or (k) an aryloxy group which may be substituted (e.g., a phenoxy or naphthoxy group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc. such as phenoxy, p-nitrophenoxy, naphthoxy etc.).

As R<sup>1</sup>, the group capable of forming an anion may have, in addition to the protective group such as an optionally substituted lower (C<sub>1-4</sub>) alkyl group or an acyl group (e.g., lower (C<sub>2-5</sub>) alkanoyl, benzoyl etc.), substituent(s) such as a lower (C<sub>1-4</sub>) alkyl group which may be substituted (the same as the "lower (C<sub>1-4</sub>) alkyl group which may be substituted" exemplified with respect to the above protective group for the group capable of forming an anion represented by R<sup>1</sup>), a halogen atom, nitro, cyano, and amino which may be substituted with one or two lower (C<sub>1-4</sub>) alkyl groups, at replaceable positions.

As R<sup>1</sup> in the above-mentioned formula, the group

capable of being converted into the group capable of forming an anion (the group having a hydrogen atom releasable as a proton) may also be a group which can be converted into the group capable of forming an anion under biological, that is, physiological conditions (e.g., reactions in the living body, such as oxidation, reduction or hydrolysis by enzymes in the living body) (so-called a prodrug), or it also may be a group such as (1) a carboxyl group, (2) a tetrazolyl group, (3) a trifluoromethanesulfonic acid amide group ( $-\text{NHSO}_2\text{CF}_3$ ), (4) a phosphate group, (5) a sulfonate group, (6) a 5- to 7-membered (preferably 5- to 6-membered) monocyclic, optionally substituted heterocyclic residue containing at least one atom selected from N, S and O, which are protected with cyano, a N-hydroxycarbamimidoyl group ( $-\text{C}(=\text{N}-\text{OH})-\text{NH}_2$ ), an optionally substituted lower ( $\text{C}_{1-4}$ ) alkyl group, and an acyl group respectively, and which can be converted by chemical reaction into the group capable of forming an anion represented by  $\text{R}^1$  (so-called a synthetic intermediate).

Preferably,  $\text{R}^1$  is carboxyl which may be protected with an optionally substituted lower ( $\text{C}_{1-4}$ ) alkyl (e.g., methyl, triphenylmethyl, methoxymethyl, ethoxymethyl, p-methoxybenzyl, p-nitrobenzyl etc.) or with an acyl group (e.g., lower ( $\text{C}_{2-5}$ ) alkanoyl, benzoyl etc.); tetrazolyl or

2,5-dihydro-5-oxo-1,2,4-oxadiazole-3-yl (preferably, tetrazolyl); or cyano, N-hydroxycarbamimidoyl (preferably, cyano), among which cyano is particularly preferably used.

In the above-mentioned formula, X indicates that the phenylene group and the phenyl group are bound to each other directly or via a spacer of a chain made of 2 or less atoms (preferably a direct bond). The spacer of a chain made of 2 or less atoms may be any divalent chain and may have a side chain insofar as the number of atoms constituting the straight chain is 1 or 2. Specifically, it is lower ( $C_{1-4}$ ) alkylene having a straight chain consisting of 1 or 2 carbon atoms, -CO-, -O-, -S-, -NH-, -CO-NH-, -O-CH<sub>2</sub>-, -S-CH<sub>2</sub>-, -CH=CH-, etc.

In the above-mentioned formula, n is an integer of 1 or 2, preferably 1.

In the formula, ring A represents a benzene ring which may further have substituent(s) other than the substituent group R<sup>2</sup>, and said substituent includes, for example, (1) halogen (e.g., F, Cl, Br etc.), (2) cyano, (3) nitro, (4) lower ( $C_{1-4}$ ) alkyl which may be substituted, (5) lower ( $C_{1-4}$ ) alkoxy, (6) an amino group which may be substituted (e.g., amino, N-lower ( $C_{1-4}$ ) alkylamino (e.g., methylamino), N,N-di-lower ( $C_{1-4}$ ) alkylamino (e.g., dimethylamino etc.), N-arylamino (e.g., phenylamino etc.), alicyclic amino (e.g., morpholino, piperidino, piperazino, N-phenylpiperazino

etc.) etc.), (7) a group represented by the formula  $\text{-CO-D'}$  wherein  $\text{D'}$  represents a hydroxyl group or a lower  $(\text{C}_{1-4})$  alkoxy which may be substituted at its alkyl moiety by a hydroxyl group, a lower  $(\text{C}_{1-4})$  alkoxy, lower  $(\text{C}_{2-6})$  alkanoyloxy (e.g., acetoxy, pivaloyloxy etc.), lower  $(\text{C}_{1-6})$  alkoxycarbonyloxy (e.g., methoxycarbonyloxy, ethoxycarbonyloxy etc.) or lower  $(\text{C}_{3-6})$  cycloalkoxycarbonyloxy (e.g., cyclohexyloxycarbonyloxy etc.), or (8) tetrazolyl, a trifluoromethanesulfonic acid amido group, a phosphate group or a sulfonate group which may be protected with acyl (e.g., lower  $(\text{C}_{2-5})$  alkanoyl, benzoyl etc.) or with lower  $(\text{C}_{1-4})$  alkyl which may be substituted (the same as the "lower  $(\text{C}_{1-4})$  alkyl group which may be substituted" exemplified with respect to the protective group for the group capable of forming an anion represented by  $\text{R}^1$ ), and the like.

The benzene ring may have been substituted at replaceable positions thereof with one or two of these substituents groups. The substituent which is possessed by A in addition to the substituent represented by  $\text{R}^2$  is preferably a lower  $(\text{C}_{1-4})$  alkyl which may be substituted (e.g., lower  $(\text{C}_{1-4})$  alkyl which may be substituted with a hydroxyl group, carboxyl group, halogen etc.), halogen etc., and more preferably, ring A does not have a substituent other than the substituent represented by  $\text{R}^2$ .

In the above-mentioned formula, the group capable of forming an anion represented by  $R^2$  (that is, a group having a hydrogen atom releasable as a proton) includes, for example, (1) a carboxyl group which may be esterified or amidated, (2) a tetrazolyl group, (3) a trifluoromethanesulfonic acid amide group ( $-NHSO_2CF_3$ ), (4) a phosphate group and (5) a sulfonate group, etc. and these may have been protected with a lower alkyl group which may be substituted (the same as the "lower ( $C_{1-4}$ ) alkyl group which may be substituted" exemplified with respect to the protective group for the group capable of forming an anion represented by  $R^1$ ) or an acyl group (e.g., lower ( $C_{2-5}$ ) alkanoyl, benzoyl etc.), and may be any group capable of forming an anion or being converted into said group chemically or biologically i.e. under physiological conditions (e.g., reactions in the living body, such as oxidation, reduction or hydrolysis in enzymes in the living body).

As  $R^2$ , the carboxyl which may be esterified or amidated includes, for example, groups represented by the formula  $-CO-D$  wherein D represents (1) a hydroxyl group, (2) amino which may be substituted (e.g., amino, N-lower ( $C_{1-4}$ ) alkylamino, N,N-di-lower ( $C_{1-4}$ ) alkylamino etc.) or (3) alkoxy which may be substituted {e.g., (i) a lower ( $C_{1-6}$ ) alkoxy group which may be substituted at its alkyl



moiety by a hydroxyl group, amino which may be substituted (e.g., amino, N-lower (C<sub>1-4</sub>) alkylamino, N,N-di-lower (C<sub>1-4</sub>) alkylamino, piperidino, morpholino etc.), halogen, lower (C<sub>1-6</sub>) alkoxy, lower (C<sub>1-6</sub>) alkylthio, lower (C<sub>3-8</sub>) cycloalkoxy, or dioxolenyl which may be substituted (e.g., 5-methyl-2-oxo-1,3-dioxolene-4-yl etc.), or (ii) groups represented by the formula -O-CH(R<sup>6</sup>)-OCOR<sup>7</sup> wherein R<sup>6</sup> represents (a) hydrogen, (b) a C<sub>1-6</sub> straight or branched lower alkyl group (e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, isopentyl, neopentyl etc.), (c) a C<sub>2-6</sub> straight or branched lower alkenyl group or (d) a C<sub>3-8</sub> cycloalkyl group (e.g., cyclopentyl, cyclohexyl, cycloheptyl etc.), and R<sup>7</sup> represents (a) a C<sub>1-6</sub> straight or branched lower alkyl group (e.g., methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl, isopentyl, neopentyl etc.), (b) a C<sub>2-6</sub> straight or branched lower alkenyl group, (c) a C<sub>1-3</sub> lower alkyl substituted with a C<sub>3-8</sub> cycloalkyl group (e.g., cyclopentyl, cyclohexyl, cycloheptyl etc.) or with an aryl group which may be substituted (e.g., a phenyl or naphthyl group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) group (e.g., benzyl, p-chlorobenzyl, phenetic, cyclopentylmethyl, cyclohexylmethyl etc.), (d) a C<sub>2-3</sub> lower alkenyl group substituted with C<sub>3-8</sub> cycloalkyl or with an aryl group which may be substituted (e.g., a phenyl

or naphthyl group which may have a halogen atom, nitro,  
lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) (e.g., a group  
such as cinnamyl having an alkenyl moiety such as vinyl,  
propenyl, allyl, isopropenyl etc.), (e) an aryl group which  
5 may be substituted (e.g., a phenyl or naphthyl group such  
as phenyl, p-tolyl, naphthyl etc., which may have a halogen  
atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.),  
(f) a C<sub>1-6</sub> straight or branched lower alkoxy group (e.g.,  
methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy,  
10 sec-butoxy, t-butoxy, n-pentyloxy, isopentyloxy,  
neopentyloxy etc.), (g) a C<sub>2-8</sub> straight or branched lower  
alkenyloxy group (e.g., allyloxy, isobutenyloxy etc.), (h)  
a C<sub>3-8</sub> cycloalkyloxy group (e.g., cyclopentyloxy,  
cyclohexyloxy, cycloheptyloxy etc.), (i) a C<sub>1-3</sub> lower alkoxy  
15 group substituted with a C<sub>3-8</sub> cycloalkyl (e.g., cyclopentyl,  
cyclohexyl, cycloheptyl etc.) or with an aryl group which  
may be substituted (e.g., a phenyl or naphthyl group which  
may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower  
(C<sub>1-4</sub>) alkoxy etc.) (e.g., a group such as benzyloxy,  
20 phenethyloxy, cyclopentylmethoxy, cyclohexylmethoxy etc.  
having an alkoxy moiety such as methoxy, ethoxy, n-propoxy,  
isopropoxy etc.), (j) a C<sub>2-3</sub> lower alkenyloxy group  
substituted with a C<sub>3-8</sub> cycloalkyl (e.g., cyclopentyl,  
cyclohexyl, cycloheptyl etc.) or with an aryl group which  
25 may be substituted (e.g., a phenyl or naphthyl group which

20251018 10:13:53

may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc.) (e.g., a group such as cinnamyloxy etc. having an alkenyloxy moiety such as vinyloxy, propenyloxy, allyloxy, isopropenyloxy etc.), or (k) an aryloxy group which may be substituted (e.g., a phenoxy or naphthoxy group which may have a halogen atom, nitro, lower (C<sub>1-4</sub>) alkyl, lower (C<sub>1-4</sub>) alkoxy etc. such as phenoxy, p-nitrophenoxy, naphthoxy etc.).

R<sup>2</sup> is preferably carboxyl which may be esterified, and examples thereof include -COOH and a salt thereof, -COOMe, -COOEt, -COOtBu, -COOPr, pivaloyloxymethoxycarbonyl, 1-(cyclohexyloxycarbonyloxy) ethoxycarbonyl, 5-methyl-2-oxo-1,3-dioxolene-4-ylmethoxycarbonyl, acetoxymethoxycarbonyl, propionyloxymethoxycarbonyl, n-butyryloxymethoxycarbonyl, isobutyryloxymethoxycarbonyl, 1-(ethoxycarbonyloxy) ethoxycarbonyl, 1-(acetoxy) ethoxycarbonyl, 1-(isobutyryloxy) ethoxycarbonyl, cyclohexylcarbonyloxymethoxycarbonyl, benzoyloxymethoxycarbonyl, cinnamyloxycarbonyl, cyclopentylcarbonyloxymethoxycarbonyl etc., and R<sup>2</sup> may be any group capable of forming an anion (e.g., COO<sup>-</sup>, derivatives thereof, etc.) or being converted into said group chemically or biologically i.e. under physiological conditions (e.g., reactions in the living body, such as oxidation, reduction or hydrolysis by enzymes in the living

body) or may be a carboxyl group or a prodrug thereof.

$R^2$  is preferably a group represented by the formula -  
CO-D wherein D represents (1) a hydroxyl group or (2) a  
lower ( $C_{1-4}$ ) alkoxy which may be substituted at its alkyl  
5 moiety by a hydroxyl group, amino, halogen, lower ( $C_{2-6}$ )  
alkanoyloxy (e.g., acetoxy, pivaloyloxy etc.), lower ( $C_{3-8}$ )  
cycloalkanoyloxy, lower ( $C_{1-6}$ ) alkoxycarbonyloxy (e.g.,  
methoxycarbonyloxy, ethoxycarbonyloxy etc.), lower ( $C_{3-8}$ )  
cycloalkoxycarbonyloxy (e.g., cyclohexyloxycarbonyloxy  
10 etc.), lower ( $C_{1-4}$ ) alkoxy or lower ( $C_{3-8}$ ) cycloalkoxy, and  
carboxyl esterified with lower ( $C_{1-4}$ ) alkyl (preferably  
methyl or ethyl) is preferable.

In the above-mentioned formula, the "hydrocarbon  
residue" in the "hydrocarbon residue which may be bound via  
15 a heteroatom and may have a substituent group" represented  
by  $R^3$  includes e.g. (1) an alkyl group, (2) an alkenyl  
group, (3) an alkynyl group, (4) a cycloalkyl group, (5) an  
aryl group, (6) an aralkyl group and the like, among which  
an alkyl group, an alkenyl group and a cycloalkyl group are  
20 preferable.

The alkyl group in the above-mentioned (1) is a lower  
alkyl group containing about 1 to 8 carbon atoms which may  
be straight or branched, and examples thereof include  
methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-  
25 butyl, t-butyl, pentyl, i-pentyl, hexyl, heptyl and octyl.

The alkenyl group in the above-mentioned (2) is a straight or branched lower alkenyl group containing about 2 to 8 carbon atoms, and examples thereof include vinyl, propenyl, 2-butenyl, 3-butenyl, isobutenyl, 2-octenyl etc.

5        The alkynyl group in the above-mentioned (3) is a lower alkynyl group containing about 2 to 8 carbon atoms which may be straight or branched, and examples thereof include ethynyl, 2-propynyl, 2-butyne, 2-pentyne, 2-octynyl etc.

10       The cycloalkyl group in the above-mentioned (4) includes a lower cycloalkyl containing about 3 to 6 carbon atoms, and examples thereof include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl etc.

15       The above-mentioned alkyl group, alkenyl group, alkynyl group or cycloalkyl group may have been substituted with a hydroxyl group, an amino group which may be substituted (e.g., amino, N-lower ( $C_{1-4}$ ) alkylamino, N,N-di-lower ( $C_{1-4}$ ) alkylamino etc.), halogen, a lower ( $C_{1-4}$ ) alkoxy group, a lower ( $C_{1-4}$ ) alkylthio group etc.

20       The aralkyl group in the above-mentioned (5) includes e.g. phenyl-lower ( $C_{1-4}$ ) alkyl such as benzyl and phenethyl, and the aryl group in the above-mentioned (6) includes e.g. phenyl.

25       The aralkyl group or the aryl group described above may have e.g. halogen (e.g., F, Cl, Br etc.), nitro, an

amino group which may be substituted (e.g., amino, N-lower  
 (C<sub>1-4</sub>) alkylamino, N,N-di-lower (C<sub>1-4</sub>) alkylamino etc.),  
 lower (C<sub>1-4</sub>) alkoxy (e.g., methoxy, ethoxy etc.), lower (C<sub>1-4</sub>)  
 alkylthio (e.g., methylthio, ethylthio etc.) or lower  
 5 (C<sub>1-4</sub>) alkyl (e.g., methyl, ethyl etc.) at an arbitrary  
 position on the benzene ring thereof.


Among the groups described above, the "hydrocarbon  
 residue" in the "hydrocarbon residue which may be bound via  
 a heteroatom and may have a substituent group" represented  
 10 by R<sup>3</sup> is preferably an alkyl or alkenyl group which may be  
 substituted (e.g., a lower (C<sub>1-5</sub>) alkyl or lower (C<sub>2-5</sub>)  
 alkenyl group which may be substituted with a hydroxyl  
 group, an amino group, halogen or a lower (C<sub>1-4</sub>) alkoxy  
 group), among which a lower (C<sub>1-5</sub>) alkyl (more preferably  
 15 ethyl) is preferable.

The "heteroatom" in the "hydrocarbon residue which may  
 be bound via a heteroatom and may have a substituent group"  
 represented by R<sup>3</sup> includes -O-, -S(O)<sub>m</sub>- [m is an integer of  
 0 to 2], and -NR'- [R' is a hydrogen atom or lower (C<sub>1-4</sub>)  
 20 alkyl], among which -O- is preferably used.


Among those described above, R<sup>3</sup> may be bound via -O-,  
 -S(O)<sub>m</sub>- [m is an integer of 0 to 2] or -NR'- [R' is a  
 hydrogen atom or lower (C<sub>1-4</sub>) alkyl], and is preferably a  
 lower (C<sub>1-5</sub>) alkyl or lower (C<sub>2-5</sub>) alkenyl group which may be  
 25 substituted with a substituent group selected from a

hydroxyl group, an amino group, halogen and a lower ( $C_{1-4}$ ) alkoxy group, particularly preferably a lower ( $C_{1-5}$ ) alkyl or lower ( $C_{1-5}$ ) alkoxy, more preferably ethoxy.

Those compounds having an angiotensin II antagonistic activity represented by the formula (I) are preferably benzimidazole-7-carboxylic acid derivative or a pharmacologically acceptable salt thereof, represented by the formula (I'):



wherein  $R^1$  represents (1) a carboxyl group, (2) a tetrazolyl group or (3) a group represented by the formula:



wherein  $i$  represents  $-O-$  or  $-S-$ ,  $j$  represents  $\geq O$ ,  $\geq S$  or  $\geq S(O)_m$  wherein  $m$  is as defined above, and ring A

represents a lower ( $C_{1-4}$ ) alkyl which may be substituted with a group besides the substituent group  $R^2$  (e.g., lower

(C<sub>1-4</sub>) alkyl which may be substituted with a hydroxyl group, a carboxyl group, halogen etc.) or a benzene ring which may be substituted with halogen (preferably a benzene ring having no substituent for the substituent represented by R<sup>2</sup>), and R<sup>2</sup> represents the formula -CO-D wherein D represents (1) a hydroxyl group or (2) a lower (C<sub>1-4</sub>) alkoxy which may be substituted at its alkyl moiety by a hydroxyl group, amino, halogen, lower (C<sub>2-6</sub>) alkanoyloxy (e.g., acetoxy, pivaloyloxy etc.), lower (C<sub>3-8</sub>) cycloalkanoyloxy, lower (C<sub>1-6</sub>) alkoxycarbonyloxy (e.g., methoxycarbonyloxy, ethoxycarbonyloxy etc.), lower (C<sub>3-8</sub>) cycloalkoxycarbonyloxy (e.g., cyclohexyloxycarbonyloxy etc.), lower (C<sub>1-4</sub>) alkoxy or lower (C<sub>3-8</sub>) cycloalkoxy, and R<sup>3</sup> may be bound via -O-, -S(O)<sub>m</sub>- [m is an integer of 0 to 2] or -NR'- [R' is a hydrogen atom or lower (C<sub>1-4</sub>) alkyl], and represents a lower (C<sub>1-5</sub>) alkyl or lower (C<sub>2-5</sub>) alkenyl group which may be substituted with a substituent selected from a hydroxyl group, an amino group, halogen and a lower (C<sub>1-4</sub>) alkoxy group (preferably lower (C<sub>1-5</sub>) alkyl or lower (C<sub>1-5</sub>) alkoxy, more preferably ethoxy), and particularly preferable used are 2-ethoxy-1-[[2'-(1H-tetrazole-5-yl)biphenyl-4-yl]methyl]benzimidazol-7-carboxylic acid [Candesartan], 1-(cyclohexyloxycarbonyloxy)ethyl 2-ethoxy-1-[[2'-(1H-tetrazole-5-yl)biphenyl-4-yl]methyl]benzimidazole-7-carboxylate [Candesartan cilexetil], pivaloyloxymethyl 2-

20251013 01:00



ethoxy-1-[[2'-(1H-tetrazole-5-yl)biphenyl-4-yl]methyl]benzimidazole-7-carboxylate, 2-ethoxy-1-[[2'-(2,5-dihydro-5-oxo-1,2,4-oxadiazole-3-yl)biphenyl-4-yl]methyl] benzimidazole-7-carboxylic acid or a salt thereof.

The above-described benzimidazole derivatives can be synthesized by a known method described in, for example, EP-425921, EP-459136, EP-553879, EP-578125, EP-520423, EP-668272 etc. or by an analogous method. When Candesartan cilexetil is used, stable C-type crystals described in EP-459136 are preferably used.

The compound having an angiotensin II antagonistic activity or a prodrug thereof used in the present invention may be used as it is or as a pharmacologically acceptable salt thereof. When the compound having an angiotensin II antagonistic activity has an acidic group such as a carboxyl group etc., the salt thereof includes those salts with inorganic bases (e.g., alkali metals such as sodium, potassium etc., alkaline earth metals such as calcium, magnesium etc., transition metals such as zinc, iron, copper etc.) and organic bases (e.g., organic amines such as trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N'-dibenzyl ethylene diamine etc., basic amino acids such as arginine, lysine, ornithine etc.).

When the compound having an angiotensin II antagonistic activity has a basic group such as an amino group etc., the salt thereof includes those salts with inorganic acids and organic acids (e.g., hydrochloric acid, nitric acid, sulfuric acid, phosphoric acid, carbonic acid, bicarbonic acid, formic acid, acetic acid, propionic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid etc.), acidic amino acids such as aspartic acid, glutamic acid etc., or the like.

In the present invention, a prodrug of the compound having an angiotensin II antagonistic activity [hereinafter, sometimes, referred to as AII antagonist] refers to a compound to be converted into an AII antagonist by reaction with enzymes, gastric acid etc. under physiological conditions in the living body, that is, a compound to be converted into an AII antagonist by enzymatic oxidation, reduction or hydrolysis or a compound to be converted into an AII antagonist by hydrolysis with gastric acid etc. A prodrug of an AII antagonist includes compounds derived from an AII antagonist by acylation, alkylation or phosphorylation of an amino group thereof (e.g., compounds wherein an amino group of an AII antagonist has been converted into eicosanoyl, alanyl, pentylaminocarbonyl, (5-

10031398-014802

methyl-2-oxo-1,3-dioxolene-4-yl)methoxycarbonyl,  
tetrahydrofuranyl, pyrrolidylmethyl, pivaloyloxymethyl,  
tert-butyl etc.); compounds derived from an AII antagonist  
by acylation, alkylation, phosphorylation or boration of a  
5 hydroxyl group thereof (e.g., compounds wherein a hydroxyl  
group of an AII antagonist has been converted into acetyl,  
palmitoyl, propanoyl, pivaloyl, succinyl, furamallyl,  
alanyl, dimethylaminomethylcarbonyl etc.); and compounds  
derived from an AII antagonist by esterifying or amidating  
10 a carboxyl group thereof (e.g., compounds wherein a  
carboxyl group of an AII antagonist has been converted into  
ethyl ester, phenyl ester, carboxymethyl ester, dimethyl  
aminomethyl ester, pivaloyloxy methyl ester,  
ethoxycarbonyloxy ethyl ester, phthalidyl ester, (5-methyl-  
15 2-oxo-1,3-dioxolene-4-yl)methyl ester, cyclohexyloxy  
carbonyloxy ethyl ester, methyl amide etc.). These  
compounds can be produced from an AII antagonist by a per  
se known method.

20 A prodrug of an AII antagonist may be the one to be  
converted into an AII antagonist under physiological  
conditions as described in "Iyakuhin no Kaihatsu"  
(Development of Pharmaceutical Preparations), vol. 7,  
"Bunshi Sekkei" (Molecular Design), pp. 163-198, published  
in 1990 by Hirokawa Shoten.

25 An AII antagonist may be a hydride or anhydride.

The compound represented by the formula (I) or a pharmaceutically acceptable salt thereof is low-toxic and can be used as an agent for preventing recurrence of cerebrovascular disorder or as an agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof in mammals (e.g., humans, mice, rats, rabbits, dogs, cats, cattle, pigs, monkeys etc.) as it is or as a pharmaceutical composition thereof with a pharmaceutically acceptable carrier.

The pharmaceutically acceptable carriers used in the present invention are conventional organic or inorganic carrier materials in pharmaceutical preparations, and are formulated as excipients, lubricants, binders or disintegrators in solid preparations, or as solvents, solubilizers, suspending agents, isotonizing agents, buffers and analgesics in liquid preparations. If necessary, pharmaceutical additives such as preservatives, antioxidants, coloring agents and sweeteners can also be used.

Preferable examples of the excipients include lactose, sucrose, D-mannitol, D-sorbitol, starch,  $\alpha$ -starch, dextrin, crystalline cellulose, low-substituted hydroxypropyl cellulose, sodium carboxymethyl cellulose, gum arabic, dextrin, pullulan, light silicic anhydride, synthetic aluminum silicate, magnesium metasilicate aluminate etc.

Preferable examples of the lubricants include magnesium stearate, calcium stearate, talc and colloidal silica.

Preferable examples of the binders include  
5   pregelatinized starch, sucrose, gelatin, gum arabic, methyl  
cellulose, carboxymethyl cellulose, sodium carboxymethyl  
cellulose, crystalline cellulose, sucrose, D-mannitol,  
trehalose, dextrin, pullulan, hydroxypropyl cellulose,  
hydroxypropylmethyl cellulose, polyvinyl pyrrolidone etc.

10       Preferable examples of the disintegrators include  
lactose, white sugar, starch, carboxymethyl cellulose,  
calcium carboxymethyl cellulose, sodium croscarmellose,  
sodium carboxymethyl starch, light silicic anhydride, low-  
substituted hydroxypropyl cellulose etc.

15       Preferable examples of the solvents include injectable  
water, physiological saline, Ringer's solution, alcohol,  
propylene glycol, polyethylene glycol, sesame oil, corn oil,  
olive oil, cottonseed oil etc.

20       Preferable examples of the solubilizers include  
polyethylene glycol, propylene glycol, D-mannitol,  
trehalose, benzyl benzoate, ethanol, trisaminomethane,  
cholesterol, triethanolamine, sodium carbonate, sodium  
citrate, sodium salicylate, sodium acetate etc.

25       Preferable examples of the suspending agents include  
surfactants such as stearyl triethanolamine, sodium

20250313 01:03:03

laurylsulfate, lauryl aminopropionic acid, lecithin, benzalconium chloride, benzetonium chloride, and glycerine monostearate; hydrophilic polymers such as polyvinyl alcohol, polyvinyl pyrrolidone, sodium carboxymethyl cellulose, methyl cellulose, hydroxymethyl cellulose, hydroxyethyl cellulose, and hydroxypropyl cellulose; Polysorbate; and polyoxyethylene hardened sesame oil.

Examples of the isotonizing agents include sodium chloride, glycerine, D-mannitol, D-sorbitol, glucose etc.

Preferable examples of the buffers include buffers such as phosphates, acetates, carbonates and citrates.

Preferable examples of the analgesics include benzyl alcohol etc.

Preferable examples of the preservatives include paraoxybenzoates, chlorobutanol, benzyl alcohol, phenethyl alcohol, hydroacetic acid, and sorbic acid. Preferable examples of the antioxidants include sulfites, ascorbates etc.

Preferable examples of the coloring agents include water-soluble edible tar pigment (e.g., edible pigments such as Edible Red Nos. 2 and 3, Edible Yellow Nos. 4 and 5, Edible Blue Nos. 1 and 2), water-insoluble lake pigments (e.g., aluminum salts of the water-soluble edible tar pigments described above), natural pigments (e.g.,  $\beta$ -carotene, chlorophyll, red iron oxide etc.) or the like.

Preferable examples of the sweeteners include sodium saccharine, glytylrytin [phonetic] dipotassium, aspartame, stevia etc.

20870-01803  
5 The dosage form of the pharmaceutical composition includes e.g. oral preparations such as tablets, capsules (including soft capsules, microcapsules etc.), granules, powders, syrups, emulsions and suspensions; and parenteral preparations such as injectable preparations (e.g., those for subcutaneous injection, intravenous injection, 10 intramuscular injection, intraperitoneal injection, injection into the glass body etc.), drips, preparations for external application (e.g., preparations for administration via the nose, transdermal preparations, ointments etc.), suppositories (e.g., rectal suppositories, 15 vaginal suppositories), pellets and drips, and these can be safely administered orally or parenterally.

The pharmaceutical preparation can be produced by conventional methods in the field of pharmaceutical manufacturing, for example by methods described in the 20 Japanese Pharmacopoeia, etc. Hereinafter, a specific process for producing the pharmaceutical preparation is described in more detail.

For example, the oral preparation can be produced by adding e.g. excipients (e.g., lactose, sucrose, starch, D- 25 mannitol etc.), disintegrators (e.g., calcium carboxymethyl

cellulose etc.), binders (e.g., pregelatinized starch, gum arabic, carboxymethyl cellulose, hydroxypropyl cellulose, polyvinyl pyrrolidone etc.) or lubricants (e.g., talc, magnesium stearate, polyethylene glycol 6000 etc.) to the active ingredient and then molding by compression, if necessary followed by coating thereof in a per se known method with a coating base with the aim of masking tastes or preparing enteric or lasting preparations.

The coating base includes e.g. a sugar coating base, a water-soluble film coating base, an enteric film coating base, a sustained release film coating base etc.

Sucrose is used as the sugar coating base, and at least one member selected from talc, precipitated calcium carbonate, gelatin, gum arabic, pullulan and carnauba wax may further be used alone or in combination.

The water-soluble film coating base includes e.g. cellulose type polymers such as hydroxypropyl cellulose, hydroxypropylmethyl cellulose, hydroxyethyl cellulose and methylhydroxyethyl cellulose; synthetic polymers such as polyvinyl acetal diethylaminoacetate, aminoalkylmethacrylate copolymer E [Eudragit E (trade name), Rhoem Pharma Ltd.], polyvinyl pyrrolidone etc.; and polysaccharides such as pullulan.

The enteric film coating base includes e.g. cellulose type polymers such as hydroxypropylmethyl cellulose

10031398-011802



phthalate, hydroxypropylmethyl cellulose acetatesuccinate, carboxymethylethyl cellulose and acetic acid phthalic acid cellulose; acrylate type polymers such as methacrylate copolymer L [Eudragit L (trade name), Rhoem Pharma Ltd.],  
 5 methacrylate copolymer LD [Eudragit L-30D55 (trade name), Rhoem Pharma Ltd.] and methacrylate copolymer S [Eudragit S (trade name), Rhoem Pharma Ltd.]; and naturally occurring materials such as shellac.

The sustained release film coating base includes e.g.  
 10 cellulose type polymers such as ethyl cellulose; aminoalkyl methacrylate copolymer RS [Eudragit RS-100 (trade name), Rhoem Pharma Ltd.] and acrylate type polymers such as an ethyl acrylate-methyl methacrylate copolymer suspension [Eudragit NE (trade name), Rhoem Pharma Ltd.].

15 Two or more of the coating bases described above may be mixed and used in a suitable ratio. For coating, e.g. sunproofing agents such as titanium oxide, iron oxide etc. may also be used.

The injectable preparation is produced by dissolving,  
 20 suspending or emulsifying the active ingredient along with a dispersant (e.g., Polysorbate 80, polyoxyethylene hardened sesame oil 60 etc.), polyethylene glycol, carboxymethyl cellulose, sodium alginate etc.), a preservative (e.g., methyl parabene, propyl parabene,  
 25 benzyl alcohol, chlorobutanol, phenol etc.), an isotonizing

2025031308 041802

agent (e.g., sodium chloride, glycerine, D-mannitol, D-sorbitol, glucose etc.) etc. in an aqueous solvent (e.g., distilled water, physiological saline, Ringer's solution etc.) or an oil solvent (e.g., propylene glycol and vegetable oils such as olive oil, sesame oil, cottonseed oil, corn oil etc.) and the like. In this case, it is possible to use additives such as a solubilizer (e.g., sodium salicylate, sodium acetate etc.), a stabilizer (e.g., human serum albumin etc.), an analgesic (e.g., benzyl alcohol etc.) or the like.

The dose of the compound represented by the formula (I) or a pharmaceutically acceptable salt thereof is varied depending on the subject of administration, administration route, intended diseases, conditions etc., but in the case of oral administration for mammals, particularly humans (weighing 50 kg), the compound represented by the formula (I) or a pharmaceutically acceptable salt thereof as the active ingredient is usually administered in a dose of about 0.001 to 500 mg, preferably 0.1 to 50 mg, and this dose is administered preferably once to thrice every day.

The agent for preventing recurrence of cerebrovascular disorder or the agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof according to the present invention can be used simultaneously or at intervals with an anti-dementia agent,

10031398-011802  
a nitrogen monoxide action enhancer, a glutamate inhibitor,  
and an inhibitor of blood vessel thickening, in addition to  
an agent for ameliorating cerebral circulation and blood  
flow, an agent for ameliorating cerebral metabolism, an  
5 agent for ameliorating mental diseases, an anti-convulsion  
agent, a therapeutic agent for hypertension, a therapeutic  
agent for diabetes, an anti-cerebral edema agent, a  
thrombolytic, an antiplatelet, an anticoagulant, an agent  
for ameliorating lipid metabolism, and a radical scavenger.

10 When these chemicals are used in combination, the  
respective chemicals can be mixed separately or  
simultaneously with pharmaceutically acceptable carriers,  
excipients, binders, diluents etc. and administered orally  
or parenterally as a pharmaceutical composition. When the  
15 chemicals are formed separately into preparations, the  
separately formed preparations can be mixed in a diluent  
and administered, or the separately formed preparations may  
be administered simultaneously or at intervals into the  
subject. The pharmaceutical preparation of the present  
20 invention also encompasses a kit product for mixing the  
separately formed preparations with a diluent just before  
use and administering them (e.g., an injection kit  
comprising ampoules containing each powdery chemical and a  
diluent for mixing and dissolving two or more chemicals)  
25 and a kit product for administering the separately formed

preparations simultaneously or at intervals to an intended subject (e.g., a kit for administering two or more kinds of tablets simultaneously or separately at intervals, comprising chemical-containing tablets introduced into an identical or different bag provided if necessary with a column thereon for describing the time of administering the chemical).

For example, the agent for preventing recurrence of cerebrovascular disorder or the agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof according to the present invention can be used in combination with agents for ameliorating cerebral circulation and blood flow, such as vinpocetine and fulnaridine, agents for ameliorating cerebral metabolism, such as anilacetam and nicergoline, agents for ameliorating mental diseases, for example therapeutic agents for schizophrenia (orazapin, risperidone etc.), anti-anxiety agents (alprazolam, diazepam etc.), anti-depression agents (fluoxcetine, certraline, fluboxamine etc.), anti-convulsion agents such as diazepam, therapeutic agents for hypertension, such as derapuryl hydrochloride and manidipine hydrochloride, therapeutic agents for diabetes, such as boguribose, pyrogritazone, sulfonyl urea agents, anti-cerebral edema agents such as glycerol, thrombolytics such as tissue plasminogen activator and prourokinase,

anti-platelet agents such as ticlopidine, clopidogrel,  
cirostazole and aspirin, anti-coagulants such as heparin,  
wafarin and argatroban, agents for ameliorating lipid  
metabolism, such as seribastatin, pravastatin and  
5      clofibrate, and radical scavengers such as vitamins E and C.

#### Best Mode for Carrying Out the Invention

Hereinafter, the present invention is described in  
more detail by reference to Examples, which however are not  
10      intended to limit the present invention.

#### Examples

The agent of the present invention for preventing  
recurrence of cerebrovascular disorder or the agent for  
ameliorating troubles following cerebrovascular disorder  
15      and inhibiting progress thereof which comprises as an  
active component the compound having an AII antagonistic  
activity or a salt thereof can be produced, for example, by  
the following formulation.

20

#### Example 1. Capsules

(1) Candesartan cilexetil	30 mg
(2) Lactose	90 mg
(3) Crystalline cellulose	70 mg
25      (4) Magnesium stearate	10 mg

10034398.011802

1 capsule

200 mg

(1), (2), (3) and 1/2 of (4) are mixed, kneaded and granulated. The remainder of (4) is added thereto, and the whole is encapsulated in a gelatin capsule.

## 5 Example 2. Tablets

(1) Candesartan cilexetil 30 mg

(2) Lactose 35 mg

(3) Corn starch 150 mg

(4) Crystalline cellulose 30 mg

10 (5) Magnesium stearate 5 mg

1 tablet 250 mg

(1), (2), (3), 2/3 of (4) and 1/2 of (5) are mixed, kneaded and granulated. The remainders of (4) and (5) are added to the granules which are then molded into a table by compression.

15

## Experimental Example 1

Action of Candesartan cilexetil in ameliorating troubles following cerebrovascular disorder in spontaneous hypertension rat with ease spontaneous cerebral apoplexy

20 (SHRSP)

## Method:

Male SHRSP are used. SHRSP are separately raised and given 1 % saline solution as drinking water to promote and regulate occurrence of cerebral apoplexy. To regulate apoplexy symptoms, rats expressing the non-voluntary

25

20031398.011802

Sub  
A2

motility of raising the foreleg are divided one after another into a control group and a drug administration group. After the first fit, the drinking water is replaced by tap water, and then nerve symptoms are observed. During  
5 observation, the rats are examined for severity of nerve symptoms and for change in their weight. After the final administration, the rats are allowed to bleed to death under anesthesia and subjected to histological investigation.

10

#### Industrial Applicability

The renin-angiotensin system is revealed to play an important role in the progress of cerebrovascular lesions accompanying hypertension. Then, a compound having an AII  
15 antagonistic activity (particularly Candesartan cilexetil) can be expected to inhibit the progress of lesions (carotid artery lesions) responsible for cerebral arteriosclerosis and ischemic cerebrovascular disorder not only by the vasodilator action but also by the action of improving  
20 endothelial cell functions and inhibiting inner membrane thickening (correction of vascular remodeling). Further, reduction in cerebral blood flow and microcirculation in brain are ameliorated by the action of ameliorating the ability for automatic circulation of cerebral circulation,  
25 and there are further brought about various actions for

20210303

Cap  
03

23  
5 protection of brain (nerves), correction of an abnormality in fibrinolytic system, amelioration of blood properties, etc. Thus, the compound is useful for preventing recurrence of cerebrovascular disorder or for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof.

10 That is, Candesartan cilexetil having the following actions can be used effectively as an agent for preventing recurrence of cerebrovascular disorder or as an agent for ameliorating troubles following cerebrovascular disorder and inhibiting progress thereof.

(1) Long-lasting action of reducing blood pressure without adversely influencing cerebral blood flow

20 It is reported that Candesartan, i.e. an active metabolite of Candesartan cilexetil, can deviate the lower limit of automatic regulation leftward (toward lower blood pressure level) (Vraamark T et al., J Hypertens 13:755-761, 1995), and in a patient with cerebrovascular disorder where the ability for automatic regulation of cerebral vessels is hindered, the recurrence of cerebrovascular disorder can be prevented by preventing or decreasing cerebral ischemia by the mechanism of blood flow kinetics as a cause of the recurrence. Further, Candesartan cilexetil can prevent recurrence of cerebrovascular disorder by its stable blood pressure depressing action through 24 hours, thus stably

25



controlling blood pressure whereby a significant reduction in blood pressure at night or an increase in blood pressure from night to early morning or in early morning, which is a cause of occurrence and recurrence of cerebrovascular disorder, is prevented. Further, Candesartan cilexetil can lower blood pressure safely to a therapeutically desired level in a patient with cerebrovascular disorder having various complications such as diabetes, cardiac diseases and renal diseases in addition to hypertension, thus significantly contributing to inhibition of the progress of organ disorders due to these complications without causing significant reduction in blood pressure, and finally the recurrence of cerebrovascular disorder can be prevented by reducing the dangerous factors for recurrence of cerebrovascular disorder.

(2) Anti-arteriosclerosis action:

It has been revealed that AII have not only a strong vasoconstricting action but also various actions such as proliferating action, inflammation action, oxidizing action, vascular penetrating action, etc., thus playing an important role in the progress of not only hypertension but also cerebrovascular lesions. That is, it is reported that AII thickens cells via expression of oncogenes and growth factors, thickens vascular walls by an increase in production of extracellular substrate, activates a

2025-09-01 09:00

Sub  
Out

transcriptional factor (NF- $\kappa$ B) to increase expression of a  
monocyte chemotactic factor (Hernandez-Presa M, et al.,  
Circulation 95, 1532-1541, 1997), and induces production of  
free radicals from inflammatory cells (Zafari AM, et al.,  
5 Hypertension 32, 488-495, 1998), thus significantly  
influencing various organ disorders including  
cerebrovascular disorder. Candesartan cilexetil not only  
inhibits these disorders attributable to AII thereby  
treating hypertension but also exerts actions for  
10 prevention of progress of arteriosclerosis, amelioration of  
vessel remodeling, amelioration of microcirculation,  
inhibition of edema, amelioration of endothelial cell  
functions (promotion of NO production in endothelial cells),  
and protection of cells thereby preventing the progress and  
15 recurrence of cerebrovascular disorder and ameliorating  
troubles after cerebrovascular disorder.

(3) Reduction in dangerous factors for diabetes etc.:

Candesartan cilexetil is known to improve insulin  
sensitivity clinically (Iimura O., et al., Am J Hypertens 8,  
20 353-357, 1995), and can ameliorate various disorders  
accompanying an abnormality in sugar resistance, diabetes  
and a reduction in insulin sensitivity as dangerous factors  
for recurrence of cerebrovascular disorder thereby  
preventing the progress and recurrence of cerebrovascular  
25 disorder and ameliorating troubles after cerebrovascular

10031398-011802

Sub  
Dis

disorder.

(4) Brain-protecting action:

Candesartan cilexetil has actions for anti-inflammation, anti-oxidization, anti-edema, amelioration of microcirculation and improvement of endothelial cell functions as described in the above (2), and can inhibit vasoconstriction and promotion of platelet agglutination caused by various vascular agonists such as endoserin and thromboxane induced and enhanced by AII, whereby the penumbra region can be saved by amelioration of blood flow and inhibition of cellular disorders at the acute stage of vascular infarction.

Further, it is reported that a disorder in the brain-blood barrier in an angiotensinogen knockout mouse is ameliorated by administration of an angiotensin peptide such as angiotensin IV (angiotensin 3-8) (Kakinuma Y. et al., Nat Med. 4, 1078-1080, 1998), and also that an angiotensin peptide such as AIV is related to memory and the action of increasing cerebral blood flow (Wright J W et al., Brain Res Rev 25, 96-124, 1997), and the increase in these peptides by administration of Candesartan cilexetil (decomposed products are increased by an increase in AII) can prevent the progress and recurrence of cerebrovascular disorder via the above-described actions and ameliorate troubles after cerebrovascular disorder.

(5) Influence on blood components:

It is reported that stimulation with AII causes an increase in hematocrit due to an increase in production of erythropoietin, an abnormality in the fibrinolytic system due to enhancement of the activity of plasminogen activator inhibitor type-1, and induction of expression of E-selectin in vascular endothelial cells and subsequent adherence of leukocytes (Grafe M et al., Cir Res 81, 804-811, 1997). These are inhibited by Candesartan cilexetil so that changes in blood such as coagulation of platelets, leukocytes and erythrocytes are ameliorated, while formation of thrombus is inhibited and microcirculation is ameliorated by the added or synergistic actions in improvement of endothelial cell functions (1) and inhibition of thromboxane production (4), whereby the progress and recurrence of cerebrovascular disorder can be prevented and troubles after cerebrovascular disorder can be ameliorated.

(6) Combination with other therapeutic agents:

Candesartan cilexetil can be used in combination with a thrombolytic, an antiplatelet, an anti-blood-coagulant (anticoagulant), an agent for ameliorating cerebral circulation, an agent for ameliorating cerebral metabolism and an agent for ameliorating mental diseases, and by these added or synergistic actions, further excellent effects can

be expected, and the dose of these agents and the frequency of administration thereof can be decreased thereby reducing side effects attributable to these chemicals.

10031398 041902  
20870 867007